

Neonatal distress

(February 2017)

Rationale

Neonatal distress is a relatively common occurrence. Failure to identify and appropriately manage an infant in distress in a timely manner can potentially lead to significant morbidity and mortality.

Causal Conditions

(list not exhaustive)

- Prematurity
- Pulmonary (e.g., meconium aspiration, pneumothorax)
- Decreased respiratory drive (e.g., maternal medications, asphyxia)
- Cardiovascular (e.g., anemia, congenital heart disease)
- Infection

Key Objectives

In cases of a neonatal distress, the candidate will be able to assess the need for and initiate resuscitation, identify causal and ongoing pathologies, and determine ongoing needs, including whether the infant requires level 2 or level 3 neonatal intensive care.

Enabling Objectives

In cases of neonatal distress, the candidate will

- list and interpret critical clinical findings, including
 - a. physical signs and symptoms that necessitate immediate resuscitation;
 - b. maternal and perinatal history;
 - c. physical examination findings relevant to formulating a differential diagnosis;

- list and interpret critical initial investigations targeted towards identifying an underlying cause (e.g., cord blood gas, blood glucose);
- construct an effective initial management plan, including
 - a. neonatal resuscitation;
 - b. elements of ongoing supportive care, including;
 - thermoregulation;
 - fluid and electrolyte balance;
 - sepsis management;
 - cardiorespiratory support;
 - c. appropriate communication with caregiver(s);
 - d. appropriate consultation or referral.